CALEB UNIVERSITY

COLLEGE OF HEALTH SCIENCES

DEPARTMENT OF NURSING SCIENCE

NSC 403

MEDICAL BIOSTATISTICS

First Semester • 2026/2027 Session

Dr. Oluyedun H.A., *DSA, MAW, FPCPharm, FPSN*

STATISTICAL REASONING • HEALTH DATA • NURSING DECISIONS

How to Use These Notes

Read each lecture carefully, then use the visual summaries to connect ideas, review procedures, and check your understanding. These diagrams are learning aids; always ensure that a statistical method fits the data, research question, and assumptions.

Course Learning Outcomes

- Define basic statistical terminologies.
- Explain various types of statistics and their uses.
- Differentiate descriptive, qualitative, and quantitative data.
- Utilise statistical methods to analyse, present, and interpret data.
- Communicate data-analysis results to support informed decisions.

Lecture Roadmap

Lecture 1: Introduction to Biostatistics and Basic Statistical Terminology
Lecture 2: Types of Data — Discrete/Categorical and Continuous Variables
Lecture 3: Descriptive and Quantitative Versus Qualitative Data
Lecture 4: Parameters and Statistics — Population Versus Sample
Lecture 5: Parametric and Non-Parametric Statistics
Lecture 6: Review of Descriptive Statistics — Measures of Central Tendency
Lecture 7: Review of Descriptive Statistics — Measures of Dispersion
Lecture 8: Introduction to Inferential Statistics
Lecture 9: Errors in Statistical Testing — Type I and Type II Errors
Lecture 10: Hypothesis Testing — Principles and Procedure
Lecture 11: Level of Significance and Degree of Freedom
Lecture 12: Drawing Inferences and Conclusions from Statistical Analysis
Lecture 13: Statistical Packages for Data Analysis
Lecture 14: Data Organisation and Interpretation — Frequency Distributions, Graphs, Charts, and Cumulative Curves
Lecture 15: Data Communication, Data-Informed Decision-Making, and the Nurse's Role in Data Collection and Quality

Preliminary Notes to You, the Student

Learning Outcomes

At the end of this course, the students should be able to:

- Define basic statistical terminologies;
- Explain various types of statistics and their uses;
- Differentiate between descriptive, qualitative and quantitative data);
- Utilise statistical methods to analyse, present and interpret data; and
- Communicate data analysis result based on informed decision.

I want you to approach this course with a specific expectation firmly in mind: biostatistics is not a mathematics course that happens to use health examples. It is a course about disciplined reasoning under uncertainty, using numbers as the vocabulary of that reasoning. Every clinical trial you will ever read about, every national health survey finding you will ever hear quoted, every quality-improvement report on your own future ward, depends on the concepts this course builds. I want you to leave these fifteen lectures able to read a statistical claim in a journal article or a health report with genuine, informed confidence — able to ask the right questions of it, rather than either accepting it uncritically or dismissing it as incomprehensible.

As with any rigorous course, I want to be transparent about the standard of evidence I hold myself to here. Any numerical example given in these lecture notes is an illustrative teaching figure, constructed specifically to demonstrate a calculation or a concept, and is not real health data. Any actual statistic you wish to cite for a real Nigerian or global health condition must be verified against current primary sources rather than drawn from an illustrative example in this document.

Lecture 1: Introduction to Biostatistics and Basic Statistical Terminology

VISUAL PROCESS MAP | Introduction to Biostatistics and Basic Statistical Terminology

Health question	Collect data	Summarise	Interpret	Use evidence

LEARNING OBJECTIVE Define biostatistics and its core basic terminology, and explain its relevance to nursing practice.

Biostatistics is the application of statistical methods to biological, medical, and public health data. I want you to notice, immediately, why this definition matters to you specifically as a future nurse, rather than remaining an abstract academic label: nursing practice generates data constantly — vital signs, medication administration records, patient outcomes, community health indicators — and biostatistics is the discipline that turns that raw data into genuine, usable knowledge.

Core Terminology

I want to build, in this very first lecture, a precise working vocabulary that every subsequent lecture in this course will depend upon.

NURSING APPLICATION | Introduction to Biostatistics and Basic Statistical Terminology

Biostatistics turns observations from patients and populations into evidence that can inform nursing decisions.

A variable is any characteristic that can differ in value from one observation to another — age, blood pressure, blood type, or treatment outcome are all variables. A population is the entire, complete group about which you wish to draw a conclusion — for instance, all pregnant women attending antenatal care in a given state. A sample is a

subset of that population, actually selected and measured, from which conclusions about the wider population are drawn. I want you to appreciate, immediately, why this distinction between population and sample matters so fundamentally: it is almost never practical to measure an entire population directly, and biostatistics exists, in substantial part, precisely to allow you to draw defensible conclusions about a population from a properly selected sample of it.

A datum (plural: data) is a single recorded observation or measurement. Data collection is the process of systematically gathering such observations. I want you to note, further, that a statistic is a numerical value calculated from sample data — a sample average, for instance — while a parameter, a term we return to and formalise properly later in this course, is the corresponding true value for the entire population, which is frequently unknown and must be estimated using a statistic calculated from a sample.

Why This Vocabulary Matters From Day One

I want you to treat this opening lecture's vocabulary as genuinely foundational, in the same way the definitions in any first pharmacology or anatomy lecture are foundational — every remaining lecture in this course will use these terms precisely, and any looseness in your own understanding of them now will compound into genuine confusion later, particularly once we reach inferential statistics and hypothesis testing.

Assignment and Application

a. Define biostatistics and explain its relevance to nursing practice.

b. Define variable, population, sample, datum, data collection, statistic, and parameter.

QUICK REVISION CHECK | Introduction to Biostatistics and Basic Statistical Terminology

Research question?	Data type?	Method appropriate?	Meaning in context?

c. Distinguish between a statistic and a parameter with one example of each.

d. Explain why a representative sample is essential for valid inference.

e. A nurse collects blood-pressure readings from 100 patients. Identify the variable, population, sample, and type of data.

Lecture 2: Types of Data — Discrete/Categorical and Continuous Variables

VISUAL PROCESS MAP | Types of Data — Discrete/Categorical and Continuous Variables

Categorical	Discrete counts	Continuous measures	Identify variable	Choose method

LEARNING OBJECTIVE Distinguish discrete/categorical data from continuous data, and explain why this distinction determines which statistical methods are appropriate.

I want to introduce, today, a classification of data that governs, more than perhaps any other single decision in the whole of this course, which statistical technique is actually appropriate for a given dataset.

Discrete (Categorical) Data

Discrete data, also called categorical data, consists of values that fall into distinct, separate categories, with no meaningful values existing between them. I want you to distinguish two subtypes.

NURSING APPLICATION | Types of Data — Discrete/Categorical and Continuous Variables

Classify the variable before selecting a summary, graph, or statistical test.

I. Nominal data consists of categories with no inherent order — blood type (A, B, AB, O), sex, or marital status are clear examples. There is no sense in which one category is "greater than" or "less than" another.
II. Ordinal data consists of categories with a genuine, meaningful order, but without a consistent, measurable distance between categories — a pain severity scale of mild, moderate, and severe is ordinal: severe is clearly greater than mild, but the "distance" between mild and moderate is not necessarily equal to the distance between moderate and severe.

Continuous Data

Continuous data consists of values that can, in principle, take any value within a given range, including fractional values, and possesses a genuine, consistent, measurable distance between any two points. Height, weight, blood pressure, and temperature are all continuous data. I want you to note a further useful distinction within continuous data.

I. Interval data has a consistent, measurable distance between values, but no true, meaningful zero point — temperature measured in degrees Celsius is the classic example, since zero degrees Celsius does not represent a genuine absence of temperature.
II. Ratio data has a consistent, measurable distance between values, and additionally possesses a true, meaningful zero point representing a genuine absence of the quantity being measured — weight is a clear example, since zero kilograms genuinely represents the complete absence of weight.

Why This Classification Genuinely Matters

I want you to appreciate, from this early stage of the course, that this classification is not an abstract taxonomic exercise. The type of data you are working with directly determines which descriptive summary measures are meaningful to calculate, and which statistical tests are appropriate to apply — calculating an average from nominal categorical data such as blood type would be a genuine, meaningless error, since there is no numerical quantity there to average, while the very same calculation is entirely appropriate and meaningful for continuous data such as blood pressure. I want you to develop the habit, whenever you encounter a new dataset for the remainder of this course, of first asking yourself which of these categories each variable belongs to, before considering any further analysis.

QUICK REVISION CHECK | Types of Data — Discrete/Categorical and Continuous Variables

Research question?	Data type?	Method appropriate?	Meaning in context?

Assignment and Application

a. Distinguish qualitative from quantitative data and give two examples of each.

b. Define nominal, ordinal, interval, and ratio levels of measurement.

c. Classify the following variables: blood type, pain score, temperature in degrees Celsius, and weight in kilograms.

d. Explain why calculating an average from nominal data is meaningless.

e. A dataset includes age, sex, blood pressure, and diagnosis. Identify the level of measurement for each variable.

Lecture 3: Descriptive and Quantitative Versus Qualitative Data

VISUAL PROCESS MAP | Descriptive and Quantitative Versus Qualitative Data

Qualitative	Describe categories	Quantitative	Measure / count	Present clearly

LEARNING OBJECTIVE Distinguish qualitative from quantitative data, and describe the purpose of descriptive statistics.

Building on the discrete-versus-continuous distinction, I want to introduce today a further, complementary classification, and to clarify a term you will use repeatedly for the remainder of this course: descriptive statistics.

Qualitative and Quantitative Data

Qualitative data describes a quality or characteristic that is not inherently numerical, even when it is sometimes coded using numbers for convenience of data entry or analysis — the nominal and ordinal categorical data introduced in the previous lecture are both, in this broader sense, qualitative in nature. Quantitative data, by contrast, represents a genuine numerical quantity that can be meaningfully measured and subjected to arithmetic operations such as addition and averaging — the continuous data introduced in the previous lecture, together with certain forms of discrete numerical data such as a genuine count (the number of children a woman has borne, for instance), fall into this category.

NURSING APPLICATION | Descriptive and Quantitative Versus Qualitative Data

The nature of the data determines how it should be organised, summarised, and interpreted.

I want you to notice, and to hold clearly in mind, that "qualitative versus quantitative" and "discrete versus continuous" are two related but genuinely distinct classification schemes, addressing slightly different questions about a given variable — the qualitative/quantitative distinction addresses whether the data represents a genuine numerical quantity at all, while the discrete/continuous distinction, applied specifically within quantitative data, addresses whether that quantity can take any value within a range or only distinct, separate values.

Descriptive Statistics, Introduced

Descriptive statistics refers to the branch of statistics concerned with summarising and describing the main features of a dataset, without attempting to draw any conclusion beyond that specific dataset itself. I want you to understand this limitation clearly and precisely, since it is a genuinely important conceptual boundary that we will formalise fully in the descriptive-statistics lectures later in this course: a descriptive statistic tells you something true about the specific sample you actually measured, and nothing more, by itself. Concluding something about a wider population from that sample requires the separate, additional apparatus of inferential statistics, addressed later in this course.

Why Correctly Classifying Your Data Is the First Genuine Step of Any Analysis

I want you to leave this lecture with a genuinely practical habit for your future research or data-analysis work, whether in an academic project or in professional practice. Before performing any analysis at all, correctly classify every variable in your dataset: is it qualitative or quantitative, and, if quantitative, is it discrete or continuous, and, if quantitative and continuous, is it interval or ratio? I want you to treat this classification step as the mandatory, non-negotiable first step of any statistical analysis, since every subsequent decision — which summary statistic to calculate, which chart to draw, which statistical test to apply — depends directly upon it.

Assignment and Application

QUICK REVISION CHECK | Descriptive and Quantitative Versus Qualitative Data

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Define descriptive statistics and explain their purpose.

b. Distinguish qualitative from quantitative data and describe the purpose of descriptive statistics.

c. Calculate the mean, median, and mode for a small dataset of blood-pressure readings.

d. Explain when the median is preferred to the mean.

e. A nurse summarises patient data using tables and graphs. Identify the type of statistics being used and justify its appropriateness.

Lecture 4: Parameters and Statistics — Population Versus Sample

VISUAL PROCESS MAP | Parameters and Statistics — Population Versus Sample

Population	Select sample	Calculate statistic	Estimate parameter	State uncertainty

LEARNING OBJECTIVE Define parameters and statistics precisely, distinguish them from one another, and explain the logic of estimating a parameter from a statistic.

I want to return today, and to formalise properly, a distinction I introduced briefly in the very first lecture of this course: the distinction between a parameter and a statistic.

Defining the Two Terms Precisely

A parameter is a numerical value that describes a characteristic of an entire population — the true average blood pressure of every adult in Nigeria, for instance, is a parameter. A statistic is a numerical value that describes a characteristic of a sample — the average blood pressure calculated from a specific group of 500 adults actually measured for a given study is a statistic.

NURSING APPLICATION | Parameters and Statistics — Population Versus Sample

A sample statistic is used to learn about a population parameter; sound inference depends on sampling quality.

I want you to notice, and to genuinely appreciate, the fundamental practical reality that makes this distinction so important: because measuring an entire population directly is almost always impractical, or entirely impossible, parameters are, in the great majority of real research and public health situations, genuinely unknown. What we actually possess is a statistic, calculated from a sample, which we use to estimate the corresponding, unknown population parameter.

Notation Convention

I want to introduce, briefly, a notational convention you will encounter in statistical literature and in any statistical software you use, since recognising it will help you read published research with greater confidence. Population parameters are conventionally denoted using Greek letters — the population mean is denoted by the Greek letter mu (μ), and the population standard deviation by the Greek letter sigma (σ). Sample statistics, by contrast, are conventionally denoted using Roman letters — the sample mean is denoted x-bar, and the sample standard deviation is denoted s. I want you to recognise this convention specifically so that, when you encounter it in a journal article or a statistical output, you immediately understand whether a given reported figure refers to an estimated sample statistic or an assumed or hypothesised population parameter.

The Logic of Estimation

I want you to appreciate the genuine, foundational logic that this distinction sets up for the remainder of this course. Because we can rarely measure a population parameter directly, biostatistics develops formal methods for using a sample statistic to make a defensible, appropriately qualified statement about the likely value of the corresponding population parameter — this is precisely the task of inferential statistics, which we build toward across the middle portion of this course, and it depends entirely on the sample being properly and representatively selected from the population in question. A sample that is not representative of its population — one collected only from a single unrepresentative clinic, for instance, rather than genuinely reflecting the broader population of interest — will produce a statistic that may estimate its own population parameter poorly, no matter how sophisticated the subsequent statistical analysis applied to it.

Assignment and Application

QUICK REVISION CHECK | Parameters and Statistics — Population Versus Sample

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Distinguish parametric from non-parametric statistical methods.

b. Explain the assumptions underlying parametric tests.

c. Give two examples of parametric tests and two examples of non-parametric tests.

d. A dataset is skewed and ordinal. Explain which family of tests is more appropriate and why.

e. A nurse compares pain scores before and after an intervention. The data are not normally distributed. Suggest an appropriate statistical test.

Lecture 5: Parametric and Non-Parametric Statistics

VISUAL PROCESS MAP | Parametric and Non-Parametric Statistics

Check data type	Check assumptions	Parametric method	or	Non-parametric method

LEARNING OBJECTIVE Distinguish parametric from non-parametric statistical methods and explain the assumptions underlying each.

I want to introduce today a further, genuinely important distinction, one that determines which family of statistical technique is appropriate for a given analysis, building directly on the data-type classification established earlier in this course.

Parametric Statistics

Parametric statistics refers to statistical methods that make specific assumptions about the underlying distribution of the population from which the data was sampled — most commonly, an assumption that the data follows a normal distribution, the symmetric, bell-shaped distribution you will encounter repeatedly throughout the remainder of this course. Parametric methods typically also require data that is at least interval-level, in the sense established earlier in this course, since they generally involve calculations, such as averaging, that require genuine numerical distance between values.

NURSING APPLICATION | Parametric and Non-Parametric Statistics

Choose a method that fits the data, study design, and assumptions—not simply the software available.

I want you to appreciate why parametric methods are generally preferred where their assumptions genuinely hold: they typically make more efficient use of the information contained in the data, and are correspondingly more likely to detect a genuine effect or difference where one truly exists, compared with the non-parametric alternative addressed next.

Non-Parametric Statistics

Non-parametric statistics refers to statistical methods that do not require a specific assumption about the underlying population distribution, making them a genuinely valuable and appropriate alternative in two common situations: where the data is ordinal or nominal, rather than genuinely interval or ratio in nature, and where continuous data does exist but is not reasonably assumed to follow a normal distribution — for instance, where a dataset is heavily skewed rather than symmetric.

I want you to understand the trade-off involved here clearly, since it is a genuinely important piece of applied judgement you will need to exercise in your own future research or data-analysis work: non-parametric methods are more broadly applicable, since they make fewer restrictive assumptions, but this greater flexibility generally comes at some cost in statistical power, meaning a non-parametric test may be somewhat less able to detect a genuine effect, of a given true magnitude, compared with an appropriately matched parametric test applied to data that genuinely does satisfy that parametric test's assumptions.

How to Decide Which Family to Use

I want you to leave this lecture with a clear, practical decision process. First, correctly classify your data type, using the framework established earlier in this course. Second, where your data is genuinely interval or ratio, assess whether it is reasonable to assume the data follows something reasonably close to a normal distribution — a judgement that can be supported through simple visual inspection of the data's distribution, or through formal statistical tests for normality that lie beyond the scope of this introductory course but which you should be aware exist. Where both these conditions are reasonably satisfied, a parametric method is generally appropriate; where either condition fails, a non-parametric alternative should generally be preferred.

QUICK REVISION CHECK | Parametric and Non-Parametric Statistics

Research question?	Data type?	Method appropriate?	Meaning in context?

Assignment and Application

a. Explain how biostatistics is used in nursing research, quality improvement, and public health.

b. Describe how a nurse can use statistics to interpret a research article or health report.

c. A nurse reads that a new intervention reduces hospital stay by 20%. List the statistical questions the nurse should ask before accepting the claim.

d. Explain how sample size, confidence intervals, and p-values influence the interpretation of research findings.

Lecture 6: Review of Descriptive Statistics — Measures of Central Tendency

VISUAL PROCESS MAP | Review of Descriptive Statistics — Measures of Central Tendency

Mean	Median	Mode	Compare values	Choose measure

LEARNING OBJECTIVE Define and calculate the three principal measures of central tendency, and explain when each is most appropriately used.

I want to turn today to a proper, systematic review of descriptive statistics, beginning with the measures used to summarise the "typical" or "central" value of a dataset.

The Three Measures

I want you to be able to define, calculate, and distinguish all three of the following measures.

NURSING APPLICATION | Review of Descriptive Statistics — Measures of Central Tendency

Central tendency describes the centre of a dataset; outliers and data type influence the best measure.

I. The mean is the arithmetic average, calculated by summing all values in a dataset and dividing by the number of values. It is, by a considerable margin, the most commonly used measure of central tendency for continuous, quantitative data.
II. The median is the middle value of a dataset when all values are arranged in ascending or descending order — for a dataset with an even number of values, it is the average of the two middle values. I want you to note that the median can be calculated for ordinal data as well as for continuous data, unlike the mean, which strictly requires genuine numerical, interval-level or ratio-level data.
III. The mode is the value that occurs most frequently within a dataset. I want you to note that the mode is the only one of these three measures that can be meaningfully calculated even for purely nominal, categorical data, since it requires only counting the frequency of each category, not any numerical ordering or arithmetic operation at all.

Choosing the Appropriate Measure

I want you to appreciate, with genuine practical importance, why choosing the correct measure of central tendency for a given dataset matters, rather than defaulting reflexively to the mean simply because it is the most familiar. The mean is genuinely sensitive to outliers — extreme values considerably higher or lower than the rest of the dataset — since every single value in the dataset directly contributes to its calculation. Consider a small dataset of household incomes in which one household's income is dramatically higher than all the others: the mean income will be pulled substantially upward by that single extreme value, potentially giving a genuinely misleading impression of what a "typical" household in that dataset actually earns. The median, by contrast, is considerably more resistant, or robust, to such outliers, since it depends only on the middle position of the ordered data, not on the actual magnitude of every individual value.

I want you to develop the practical habit, whenever you encounter a dataset genuinely likely to be skewed — asymmetrically distributed, with a long tail of unusually high or unusually low values — of considering the median as a more genuinely representative measure of central tendency than the mean, or, at minimum, of reporting both measures together so that a reader can judge for themselves whether the two differ meaningfully, itself a useful indicator of skew.

Assignment and Application

a. Define mean, median, and mode.

b. Calculate the mean, median, and mode for a given dataset.

QUICK REVISION CHECK | Review of Descriptive Statistics — Measures of Central Tendency

Research question?	Data type?	Method appropriate?	Meaning in context?

c. Explain when each measure is most appropriate.

d. A dataset includes an extreme outlier. Explain the effect on the mean and median.

e. A nurse summarises patient ages. Explain which measure of central tendency is most informative.

Lecture 7: Review of Descriptive Statistics — Measures of Dispersion

VISUAL PROCESS MAP | Review of Descriptive Statistics — Measures of Dispersion

Range	Variance	Standard deviation	Interquartile range	Report spread

LEARNING OBJECTIVE Define and calculate the principal measures of dispersion, and explain why central tendency alone is insufficient to describe a dataset.

I want you to begin this lecture by considering a genuinely important limitation of everything covered in the previous lecture: two datasets can share the exact same mean, and yet be genuinely, meaningfully different from one another in a way that the mean alone completely fails to capture. Measures of dispersion — also called measures of spread or variability — describe how widely the values within a dataset are scattered around their central value, and I want you to appreciate, from this opening observation, why no proper description of a dataset is ever complete using a measure of central tendency alone.

The Range

The range is the simplest measure of dispersion, calculated as the difference between the highest and lowest values in a dataset. I want you to note its principal limitation directly: because it depends only on the two most extreme values, the range is even more sensitive to outliers than the mean, and provides no information whatsoever about how the remaining values in between those two extremes are actually distributed.

NURSING APPLICATION | Review of Descriptive Statistics — Measures of Dispersion

A centre without a measure of spread can conceal important differences between datasets.

Variance and Standard Deviation

The variance measures the average of the squared deviations of each value from the dataset's mean. I want you to note why the deviations are specifically squared before being averaged: if they were not squared, the positive and negative deviations from the mean would simply cancel each other out algebraically, always summing to exactly zero regardless of how widely the data was actually spread, which would make the resulting measure entirely useless.

The standard deviation is calculated as the square root of the variance, and I want you to appreciate why this final step matters genuinely, and is not a merely cosmetic mathematical flourish: because the variance is calculated using squared deviations, its own units are the square of the original data's units — if you are measuring blood pressure in millimetres of mercury, the variance is expressed in millimetres of mercury squared, a unit with no direct, intuitive interpretation. Taking the square root returns the measure to the original, directly interpretable units of the underlying data, which is precisely why the standard deviation, rather than the variance itself, is the measure of dispersion you will encounter reported most frequently in published research and clinical data.

The Interquartile Range

The interquartile range (IQR) measures the spread of the middle 50 percent of a dataset, calculated as the difference between the value at the 75th percentile (the third quartile) and the value at the 25th percentile (the first quartile). I want you to recognise the interquartile range as the natural companion measure of dispersion to the median introduced in the previous lecture, since both share the same genuine robustness to outliers, in clear contrast to the mean-and-standard-deviation pairing, both of which are more sensitive to extreme values.

Reporting Central Tendency and Dispersion Together

QUICK REVISION CHECK | Review of Descriptive Statistics — Measures of Dispersion

Research question?	Data type?	Method appropriate?	Meaning in context?

I want you to leave this lecture with a firm, standing rule for your own future data reporting: a measure of central tendency should never be reported alone, without an accompanying measure of dispersion. Reporting a mean with its standard deviation, or a median with its interquartile range — pairing measures with matched robustness properties, as just established — together give a reader a genuinely complete initial picture of a dataset's shape, in a way that either measure reported entirely alone simply cannot provide.

Assignment and Application

a. Define range, variance, standard deviation, and interquartile range.

b. Calculate the range, variance, and standard deviation for a given dataset.

c. Explain the importance of measures of dispersion.

d. Two datasets have the same mean but different standard deviations. Explain the implications.

e. A nurse compares the variability of blood-pressure readings in two groups. Explain which measure is most appropriate.

Lecture 8: Introduction to Inferential Statistics

VISUAL PROCESS MAP | Introduction to Inferential Statistics

Sample data	Estimate	Quantify uncertainty	Test / interval	Population inference

LEARNING OBJECTIVE Define inferential statistics, explain its relationship to descriptive statistics, and introduce the concept of sampling variability.

I want to turn today to the second great branch of statistics, building directly on the population-versus-sample distinction established earlier in this course, and on the recognition that descriptive statistics, on its own, is limited to describing only the specific sample actually measured.

Defining Inferential Statistics

Inferential statistics refers to the branch of statistics concerned with drawing conclusions, and making generalisations, about a population, based on data collected from a sample of that population. I want you to appreciate the genuinely important conceptual leap this represents beyond descriptive statistics: where a descriptive statistic simply reports what was observed in a specific sample, an inferential statistical statement makes a claim that extends beyond that sample, to the wider population it was drawn from — and, correspondingly, carries with it a genuine degree of uncertainty, which the formal methods of inferential statistics are specifically designed to quantify honestly, rather than to eliminate.

NURSING APPLICATION | Introduction to Inferential Statistics

Inferential statistics use sample evidence to estimate population values or assess hypotheses, with uncertainty stated.

Sampling Variability

I want to introduce, today, a concept genuinely central to understanding why inferential statistics is necessary at all, and why it must always be accompanied by an honest quantification of uncertainty. Sampling variability refers to the fact that different samples drawn from the very same population will, purely by chance, produce somewhat different sample statistics from one another, even when the underlying population parameter itself has not changed at all. I want you to work through a concrete illustration: if you were to draw many different, independent samples of 100 patients each from the very same hospital population, and calculate the average blood pressure within each separate sample, you would very likely obtain a range of somewhat different average values across those different samples, purely due to the play of chance in exactly which 100 patients happened to be selected each time — not because the true, underlying population average blood pressure was itself actually changing between samples.

Why This Concept Justifies the Entire Remainder of This Course

I want you to appreciate, quite directly, why sampling variability is the single conceptual justification for everything we build across the remaining lectures of this course. Because any single sample statistic could, purely by chance, differ somewhat from the true population parameter it is intended to estimate, we need formal methods — hypothesis testing, significance levels, confidence intervals — that honestly quantify just how much a given sample statistic might plausibly differ from the truth, purely due to this unavoidable, inherent sampling variability, rather than presenting a single sample statistic as though it were a certain, exact statement about the entire population. This is precisely the task the remaining lectures of this course build toward, piece by piece.

Assignment and Application

a. Define probability and explain its relevance to biostatistics.

b. Describe the normal distribution and its properties.

QUICK REVISION CHECK | Introduction to Inferential Statistics

Research question?	Data type?	Method appropriate?	Meaning in context?

c. Explain the concept of z-scores and standardisation.

d. A variable is normally distributed with a mean of 100 and a standard deviation of 15. Calculate the z-score for a value of 130.

e. Explain the importance of the normal distribution in statistical inference.

Lecture 9: Errors in Statistical Testing — Type I and Type II Errors

VISUAL PROCESS MAP | Errors in Statistical Testing — Type I and Type II Errors

Reality	Test decision	Type I error	Type II error	Balance risks

LEARNING OBJECTIVE Define Type I and Type II errors, and explain the practical trade-off between them.

I want to introduce today a concept of genuine importance to correctly interpreting any statistical test result you will ever encounter, in this course or in your future professional reading: the two distinct types of error a statistical test can make.

The Logical Setup

Before defining the two error types, I want you to understand the basic logical structure every statistical hypothesis test shares. A null hypothesis proposes that there is genuinely no effect, no difference, or no association present — for instance, that a new drug produces no genuine difference in outcome compared with an existing standard treatment. An alternative hypothesis proposes that a genuine effect, difference, or association does exist. A statistical test uses sample data to make a decision about whether to reject the null hypothesis in favour of the alternative, or to retain the null hypothesis as a reasonable working position given the available evidence. Because this decision is made from sample data, rather than from complete knowledge of the entire population, this decision can, of course, sometimes be genuinely wrong.

NURSING APPLICATION | Errors in Statistical Testing — Type I and Type II Errors

Type I error is a false positive; Type II error is a false negative. Context affects their consequences.

Type I Error

A Type I error occurs when you reject a null hypothesis that is, in truth, actually correct — concluding that a genuine effect exists when, in reality, it does not. I want you to note the common shorthand for this error, since you will frequently encounter it in published literature: it is sometimes called a "false positive."

Type II Error

A Type II error occurs when you fail to reject a null hypothesis that is, in truth, actually false — concluding that no genuine effect exists when, in reality, one genuinely does. I want you to note the corresponding shorthand: it is sometimes called a "false negative."

The Genuine, Unavoidable Trade-Off

I want you to appreciate, with real practical seriousness, that these two error types exist in a genuine trade-off relationship with one another, given a fixed sample size: all else being equal, reducing the probability of one error type tends to increase the probability of the other. Making a test more conservative — genuinely harder to reject the null hypothesis — reduces the risk of a Type I error, but correspondingly increases the risk of a Type II error, since a genuine effect becomes correspondingly harder to detect and confirm statistically. I want you to connect this directly to a clinical, practical example worth genuinely internalising: in the evaluation of a new medication, a Type I error might lead to a genuinely ineffective drug being wrongly approved as effective, while a Type II error might lead to a genuinely effective, beneficial drug being wrongly rejected as ineffective — both represent genuine, real-world harms, of different but comparably serious kinds, which is precisely why researchers must think carefully and explicitly about this trade-off when designing a study, rather than focusing exclusively on minimising only one of the two error types while ignoring the other entirely.

QUICK REVISION CHECK | Errors in Statistical Testing — Type I and Type II Errors

Research question?	Data type?	Method appropriate?	Meaning in context?

Assignment and Application

a. Define sampling and explain the different sampling methods.

b. Describe the concept of sampling distribution.

c. Explain the central limit theorem and its importance.

d. A nurse selects a sample of patients for a study. Explain the importance of random sampling.

e. Explain how sample size affects the precision of estimates.

Lecture 10: Hypothesis Testing — Principles and Procedure

VISUAL PROCESS MAP | Hypothesis Testing — Principles and Procedure

State H_0 and H_1	Choose test	Set α	Calculate result	Conclude in context

LEARNING OBJECTIVE Describe the standard procedure of statistical hypothesis testing, and explain the role of the p-value within it.

Building directly on last lecture's framework, I want to walk you through today the standard, systematic procedure followed in conducting a formal statistical hypothesis test.

The Standard Procedure

I want you to be able to name and describe each of the following sequential steps.

NURSING APPLICATION | Hypothesis Testing — Principles and Procedure

A statistical conclusion should follow a clear procedure and answer the research question.

I. State the null and alternative hypotheses, precisely and clearly, before any data is analysed — a step that must genuinely occur first, in principle, to avoid the serious methodological error of adjusting your hypothesis after already having seen the data's results.
II. Choose an appropriate significance level, conventionally denoted by the Greek letter alpha (α), representing the maximum acceptable probability of a Type I error you are willing to tolerate — we formalise this concept properly in the following lecture.
III. Select and apply an appropriate statistical test, chosen according to the data type and study design involved, drawing directly on the parametric-versus-non-parametric distinction established earlier in this course.
IV. Calculate the test statistic and corresponding p-value from the sample data.
V. Compare the p-value against the chosen significance level, and make a formal decision to either reject or retain the null hypothesis on this basis.
VI. Interpret and report the result in the appropriate clinical or scientific context, a genuinely important final step we return to and develop more fully later in this course.

Understanding the P-Value

I want you to be able to define the p-value with genuine precision, because it is very frequently misunderstood, even among practising health professionals, and I want you to leave this course able to state its correct meaning with real confidence. The p-value is the probability of obtaining a result at least as extreme as the one actually observed, if the null hypothesis were, in fact, true. I want you to notice, and to actively guard against, a very common and genuinely significant misinterpretation: the p-value is not the probability that the null hypothesis itself is true, nor is it the probability that the observed result occurred purely by chance in some vague, general sense. It is a specific, conditional probability, calculated on the explicit assumption that the null hypothesis holds — and I want you to be able to correct this common misinterpretation confidently whenever you encounter it, whether in a classmate's explanation, in a piece of published writing, or in your own initial thinking as you first work through this material.

A Small Worked Illustration

Consider an illustrative example, using constructed teaching figures rather than any real dataset: a study compares the mean recovery time between two patient groups, and a statistical test yields a p-value of 0.03. If your chosen significance level is 0.05, you would reject the null hypothesis, since 0.03 is less than 0.05, concluding that the observed difference in recovery time is unlikely to have arisen purely by the play of chance alone, under the assumption that no true underlying difference genuinely exists between the two groups.

Assignment and Application

QUICK REVISION CHECK | Hypothesis Testing — Principles and Procedure

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Define point estimate and interval estimate.

b. Explain the concept of confidence interval.

c. Calculate a 95% confidence interval for a mean.

d. Explain how to interpret a confidence interval.

e. A study reports a mean difference with a 95% confidence interval. Explain the meaning.

Lecture 11: Level of Significance and Degree of Freedom

VISUAL PROCESS MAP | Level of Significance and Degree of Freedom

Choose α	Assess evidence	Degrees of freedom	Reference distribution	Report result

LEARNING OBJECTIVE Define the level of significance and degree of freedom, and explain the practical role of each in hypothesis testing.

I want to formalise, today, two further concepts already referenced in the previous two lectures, giving each the dedicated, precise treatment it genuinely deserves.

Level of Significance

The level of significance, denoted alpha (α), is the threshold probability of a Type I error that a researcher is willing to accept before conducting a statistical test, chosen and fixed in advance of analysing the data. I want you to note the conventional value most commonly adopted across health research, though I want you to understand it as a widely used convention rather than a rigid, universally mandated mathematical law: an alpha level of 0.05, corresponding to a 5 percent acceptable risk of a Type I error, is by far the most commonly used threshold in published health research.

NURSING APPLICATION | Level of Significance and Degree of Freedom

Significance level and degrees of freedom affect the interpretation of many statistical tests.

I want you to appreciate a genuinely important point of nuance here: choosing a smaller alpha level, such as 0.01, makes a test more conservative, reducing the acceptable risk of a Type I error, but, connecting directly back to the trade-off established two lectures ago, correspondingly increases the risk of a Type II error for any given, fixed sample size. Some research contexts — where a false positive finding would carry particularly serious consequences, such as approving a genuinely harmful treatment — may deliberately choose a smaller, more conservative alpha level than the conventional 0.05, precisely to reduce the risk of exactly this more serious type of error.

Degree of Freedom

Degree of freedom refers to the number of independent values, or pieces of information, genuinely free to vary when calculating a given statistic, after accounting for any constraints already imposed on the data. I want you to work through a simple, concrete illustration to build genuine intuition for this concept, rather than treating it as an abstract formula to be applied mechanically. If you know that a set of five numbers has a mean of exactly 10, then only four of those five numbers are genuinely free to take any value at all; the fifth number is automatically and entirely determined by the requirement that the full set of five values must average to exactly 10. In this specific illustration, the degree of freedom is therefore four — one less than the total sample size, reflecting the one constraint imposed by fixing the mean.

Why Degree of Freedom Genuinely Matters in Practice

I want you to appreciate why this concept is not merely a mathematical curiosity you can safely set aside once this lecture ends. Many commonly used statistical distributions used in hypothesis testing — the t-distribution being a particularly important example you will very likely encounter in further, more advanced statistical coursework — change their precise shape depending on the specific degree of freedom involved. Correctly calculating and reporting degree of freedom, alongside your chosen significance level and your calculated test statistic, is therefore a genuinely necessary, rather than merely a stylistic or decorative, part of correctly conducting and fully reporting any formal statistical test.

Assignment and Application

QUICK REVISION CHECK | Level of Significance and Degree of Freedom

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Define hypothesis, null hypothesis, and alternative hypothesis.

b. Explain the steps in hypothesis testing.

c. Define type I and type II errors.

d. Explain the concept of p-value and significance level.

e. A study reports a p-value of 0.03. Explain the meaning.

Lecture 12: Drawing Inferences and Conclusions from Statistical Analysis

VISUAL PROCESS MAP | Drawing Inferences and Conclusions from Statistical Analysis

Read result	Assess uncertainty	Check assumptions	Clinical relevance	Qualified conclusion

LEARNING OBJECTIVE Describe the principles of drawing sound, appropriately qualified conclusions from a statistical analysis, and distinguish statistical significance from clinical or practical significance.

I want to address, today, a genuinely crucial final step that a great many students, and even some published researchers, handle less carefully than they should: how to move responsibly from a completed statistical calculation to a genuinely sound, well-qualified conclusion.

Statistical Significance Versus Clinical Significance

I want you to fix a distinction firmly in mind that I consider one of the single most practically important ideas in this entire course. Statistical significance refers strictly to the specific, technical judgement that a hypothesis test's result met the chosen significance-level threshold established two lectures ago — nothing more, and nothing less than that specific technical claim. Clinical (or practical) significance, by clear contrast, refers to whether an observed effect or difference is actually large enough to matter genuinely in real clinical or practical terms, for real patients or real health systems.

NURSING APPLICATION | Drawing Inferences and Conclusions from Statistical Analysis

Statistical significance does not automatically mean clinical importance or prove causation.

I want you to work through why these two concepts can, quite genuinely, come apart from one another in practice. With a sufficiently large sample size, even a genuinely tiny, clinically trivial difference between two groups can become statistically significant, simply because a very large sample size reduces sampling variability to the point where even a very small true difference can be reliably detected by a formal test. I want you to internalise, as a firm and lasting professional habit, that a "statistically significant" result is never, by itself, automatically a clinically important or a practically meaningful one — you must always separately, explicitly ask whether the actual observed magnitude of the effect is large enough to matter genuinely for real patient care or real health policy, a judgement that statistical significance alone can never make for you.

Confidence Intervals as a Richer Alternative to a Bare P-Value

I want to introduce, briefly, a further reporting tool that directly addresses part of this concern, and which you should actively look for, and value, whenever you read published research: a confidence interval provides a plausible range of values within which the true, unknown population parameter is likely to lie, rather than the single, binary reject-or-retain decision a bare p-value alone provides. I want you to appreciate the genuine, practical advantage a confidence interval offers over a p-value reported entirely alone: it directly communicates both the direction and the plausible magnitude of an effect, information a bare p-value, by itself, simply does not convey at all.

Sound Practice in Drawing Conclusions

I want you to leave this lecture with a firm, standing set of habits for your own future analytical and research work. State your conclusion in language that is properly, honestly qualified — "the data are consistent with," or "provide evidence for," rather than language of false, unwarranted certainty. Explicitly distinguish statistical significance from clinical significance in your own reporting, every single time. And never overstate a finding beyond what your specific study design and your specific sample size can genuinely, defensibly support.

Assignment and Application

QUICK REVISION CHECK | Drawing Inferences and Conclusions from Statistical Analysis

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Describe the t-test, chi-square test, and ANOVA.

b. Explain when each test is appropriate.

c. A nurse compares the mean blood pressure of two groups. Suggest an appropriate test.

d. A nurse compares the proportion of patients with an outcome in two groups. Suggest an appropriate test.

e. Explain the assumptions of each test.

Lecture 13: Statistical Packages for Data Analysis

VISUAL PROCESS MAP | Statistical Packages for Data Analysis

Prepare dataset	Check coding	Run analysis	Verify output	Report transparently

LEARNING OBJECTIVE Describe the role and general features of statistical software packages used in health data analysis.

I want to turn today from statistical theory to statistical practice, addressing the software tools you will very likely encounter and use, whether in an academic research project during your training or in professional practice afterward.

Why Statistical Software Exists

I want you to appreciate why dedicated statistical software genuinely matters, rather than treating this lecture as merely a survey of unrelated product names. The calculations underlying even a moderately sized real dataset — means, standard deviations, hypothesis tests applied across many variables simultaneously — become genuinely impractical to perform reliably by hand once a dataset grows beyond a small illustrative example of the kind used earlier in this course. Statistical software packages exist specifically to perform these calculations reliably, consistently, and rapidly, freeing you, the researcher or analyst, to focus your own attention on correctly interpreting the results, rather than on the mechanical arithmetic of producing them.

NURSING APPLICATION | Statistical Packages for Data Analysis

Software performs calculations; the researcher remains responsible for correct method choice and interpretation.

Commonly Encountered Packages

I want you to be aware of several widely used statistical packages, without this course requiring hands-on proficiency in any specific one, since that is properly the domain of a dedicated, separate practical training session or module. SPSS (Statistical Package for the Social Sciences) is very widely used across health and nursing research specifically, valued particularly for its accessible, largely menu-driven interface that does not strictly require programming skill to operate. Other packages you may encounter in further study or in the published literature include general-purpose statistical and data-analysis software with a stronger programming orientation, offering greater flexibility for more advanced or highly customised analysis, at the cost of a somewhat steeper initial learning curve than a purely menu-driven package typically presents.

The Genuinely Important Caution I Want You to Carry From This Lecture

I want you to hold onto a single, genuinely important caution from this entire lecture, one I consider considerably more important than knowing any specific software package's name or interface. Statistical software will readily, obediently perform whatever calculation you instruct it to perform, entirely regardless of whether that specific calculation is actually appropriate for your particular data type, your particular study design, or your particular research question. A statistical package cannot, and will not, warn you that you have applied a parametric test to data that plainly violates a parametric test's basic underlying assumptions, or that you have calculated a mean for data that is, in fact, purely nominal and categorical in nature. This is precisely why everything established across the earlier lectures of this course — correctly classifying your data type, correctly choosing between parametric and non-parametric methods, correctly understanding what a p-value actually does and does not mean — remains genuinely essential conceptual knowledge, entirely independent of, and prior to, whichever specific software tool you eventually happen to use to perform your calculations.

Assignment and Application

a. Define correlation and regression.

b. Explain the difference between correlation and causation.

QUICK REVISION CHECK | Statistical Packages for Data Analysis

Research question?	Data type?	Method appropriate?	Meaning in context?

c. Calculate and interpret a correlation coefficient.

d. Describe simple linear regression and its uses.

e. A nurse examines the relationship between age and blood pressure. Explain which analysis is appropriate.

Lecture 14: Data Organisation and Interpretation — Frequency Distributions, Graphs, Charts, and Cumulative Curves

VISUAL PROCESS MAP | Data Organisation and Interpretation — Frequency Distributions, Graphs, Charts, and Cumulative Curves

Raw observations	Frequency table	Choose chart	Inspect pattern	Interpret accurately

LEARNING OBJECTIVE Describe methods for organising and visually presenting data, including frequency distribution tables, common graph and chart types, and cumulative frequency curves.

I want to turn today to the practical tools used to organise raw data into a form that can be genuinely, readily understood, both by yourself as the analyst and by whoever you are ultimately communicating your findings to.

Frequency Distribution Tables

A frequency distribution table organises raw data by listing each distinct value, or each defined range of values (a class interval), together with the number of observations, or frequency, falling into each. I want you to note a further, genuinely useful column commonly included alongside frequency: relative frequency, expressing each frequency as a proportion or percentage of the total number of observations, which allows for a meaningful, direct comparison between datasets of genuinely different total sizes.

NURSING APPLICATION | Data Organisation and Interpretation — Frequency Distributions, Graphs, Charts, and Cumulative Curves

Use a display suited to the data type and avoid scales or labels that could mislead readers.

Common Graph and Chart Types

I want you to be able to name each of the following, and to identify which is appropriate for which data type, drawing directly on the classification established earlier in this course.

I. A bar chart displays the frequency of each category in categorical (nominal or ordinal) data, using separated bars, with a genuinely meaningful, deliberate gap between each bar, reflecting the fact that the underlying categories are discrete and unconnected to one another.
II. A histogram displays the frequency distribution of continuous data, grouped into class intervals, using adjoining bars with no gap between them, reflecting the fact that the underlying continuous variable flows without interruption from one class interval directly into the next.
III. A pie chart displays the relative proportion of each category within a whole, as differently sized wedges of a single circle, and is generally most appropriate and most easily interpreted when the number of distinct categories being displayed is relatively small.
IV. A line graph displays how a variable changes over a continuous sequence, most commonly over time, and is particularly well suited to displaying a genuine trend.

I want you to note, and to actively avoid in your own future work, a genuinely common and easily made error: using a pie chart for continuous data, or a histogram for purely nominal categorical data, represents a mismatch between visualisation technique and underlying data type, of exactly the same fundamental kind as applying an inappropriate statistical test to data that does not actually suit it.

Cumulative Frequency and the Cumulative Frequency Curve

Cumulative frequency is the running, progressively accumulating total of frequencies up to and including a given class interval, calculated by successively adding each interval's own frequency to the sum of all frequencies that came before it. A cumulative frequency curve (sometimes called an ogive) plots these cumulative frequency values, and I want you to appreciate its genuinely distinctive practical use: it allows you to directly read off, from the graph

itself, the approximate number, or proportion, of observations falling below any given specific value — a task that a simple, non-cumulative histogram cannot perform nearly as directly or conveniently.

QUICK REVISION CHECK | Data Organisation and Interpretation — Frequency Distributions, Graphs, Charts, and Cumulative Curves

Research question?	Data type?	Method appropriate?	Meaning in context?

Choosing the Right Presentation Method

I want you to leave this lecture with a clear, standing principle: the purpose of any data visualisation is to make a genuine pattern in the data more immediately, intuitively clear to your reader than a raw table of numbers alone could ever achieve. Choosing the correct method for your specific data type, as outlined across this lecture, is what makes that genuine communicative purpose actually achievable, rather than the visualisation itself becoming, through mismatch or misuse, simply another potential source of confusion or misinterpretation.

Assignment and Application

a. Describe the uses of statistical software in biostatistics.

b. Explain the principles of effective data presentation.

c. Compare tables, bar charts, histograms, and line graphs.

d. A nurse prepares a report on patient outcomes. Explain how to present the data effectively.

e. Explain the importance of transparency and accuracy in data presentation.

Lecture 15: Data Communication, Data-Informed Decision-Making, and the Nurse's Role in Data Collection and Quality

VISUAL PROCESS MAP | Data Communication, Data-Informed Decision-Making, and the Nurse's Role in Data Collection and Quality

Collect accurately	Check completeness	Analyse appropriately	Communicate clearly	Decide and improve

LEARNING OBJECTIVE Describe principles of effective data communication, explain the concept of data-informed decision-making, and articulate the specific responsibilities of the nurse in data collection and data quality.

I want to close this course today by bringing together its statistical content and connecting it directly, concretely, to your own professional identity and daily responsibility as a nurse.

Principles of Effective Data Communication

I want you to be able to name several principles that genuinely distinguish effective data communication from a mere accurate recitation of numbers. Effective communication is audience-appropriate — a report intended for fellow statisticians can reasonably include considerably more technical detail than a report intended for a community health committee or a group of patients, without either audience being poorly served. It is honest about uncertainty, actively avoiding the false, unwarranted appearance of exact, perfect precision where genuine uncertainty exists, drawing directly on the confidence-interval reasoning established earlier in this course. It separates description from interpretation clearly, so that a reader can readily distinguish what the data actually showed from what the analyst concluded or recommends on the basis of that data. And it uses appropriate visualisation, drawing on the correctly matched graph and chart types established in the previous lecture, rather than either overwhelming a reader with unnecessary technical detail or, at the opposite extreme, oversimplifying a genuinely complex finding into something misleadingly straightforward.

NURSING APPLICATION | Data Communication, Data-Informed Decision-Making, and the Nurse's Role in Data Collection and Quality

Reliable decisions depend on accurate source data, honest communication, and appropriate interpretation.

Data-Informed Decision-Making

Data-informed decision-making refers to the deliberate practice of using systematically collected and properly analysed data, rather than intuition, anecdote, or habit alone, to actually guide clinical, managerial, or policy decisions. I want you to appreciate why this matters directly and concretely to you as a nurse, and not merely to researchers or to central policy-makers working at a considerable distance from the bedside: a ward-level decision about staffing patterns, informed by properly analysed patient-acuity data, or a community health decision about where to concentrate a limited immunisation outreach effort, informed by properly analysed local surveillance data, are both genuine, everyday examples of exactly the kind of data-informed decision-making this concept refers to — decisions of the kind you may find yourself directly involved in making, or contributing evidence toward, throughout your own future nursing career.

The Nurse's Role in Data Collection and Data Quality

I want to close this entire course by making this final, concrete connection as explicit and as practically grounded as I possibly can. Nurses are very frequently the health professionals who actually generate the raw data that any subsequent statistical analysis, however sophisticated, ultimately depends upon entirely — vital signs recorded on a chart, a medication administered and documented, a case reported through a surveillance system. I want you to appreciate, quite directly, that no statistical technique covered anywhere across the preceding fourteen lectures of this course, however correctly and carefully applied, can ever meaningfully compensate for poor-quality data collected carelessly at its original, initial source. A statistician working with inaccurate, incomplete, or inconsistently recorded data will, however sophisticated their subsequent analysis, produce an unreliable,

ultimately misleading result regardless of that sophistication — a principle sometimes summarised, quite memorably, by the phrase "garbage in, garbage out."

Your specific responsibilities as a nurse in relation to data quality include: recording observations accurately, precisely as observed, rather than as you might expect or assume them to be; recording observations completely, without leaving important fields blank or omitted; recording observations in a timely manner, since delayed recording genuinely increases the real risk of inaccurate recall; and using standardised definitions and standardised forms, consistently, so that data recorded by different nurses, on different shifts, at different facilities, can be meaningfully and validly combined and compared with one another during any subsequent analysis. I want you to recognise, in closing, that every single statistical concept covered across the whole of this course — from the very first lecture's basic terminology, through descriptive and inferential statistics, to today's discussion of communication and data-informed decision-making — depends, at its very root, on the quality of data you yourself, as a practising nurse, will be personally and directly responsible for generating throughout the whole of your career.

Assignment and Application

QUICK REVISION CHECK | Data Communication, Data-Informed Decision-Making, and the Nurse's Role in Data Collection and Quality

Research question?	Data type?	Method appropriate?	Meaning in context?

a. Explain the role of biostatistics in evidence-based practice.

b. Describe how biostatistics supports quality improvement in healthcare.

c. A nurse reads a systematic review and meta-analysis. Explain how to interpret the statistical findings.

d. Explain the importance of critical appraisal of statistical methods in research.

e. Reflect on how biostatistics integrates with nursing practice to improve patient care.

References and Further Reading

The following are standard, internationally recognised reference works. You should consult the current edition of each, as content is periodically revised.

Daniel, W.W. & Cross, C.L. Biostatistics: A Foundation for Analysis in the Health Sciences. Wiley.

Pagano, M. & Gauvreau, K. Principles of Biostatistics. CRC Press.

World Health Organization. Basic Epidemiology (current edition), for its treatment of health data sources and interpretation.

IBM Corporation. SPSS Statistics documentation (current edition).

Nigeria Centre for Disease Control (NCDC). Surveillance data quality and reporting guidance.